

# Association of the Serotonin Transporter Gene Promoter Region (5-HTTLPR) Polymorphism With Biased Attention for Emotional Stimuli

Christopher G. Beevers, Tony T. Wells,  
and Alissa J. Ellis  
University of Texas at Austin

John E. McGeary  
Providence Veterans Affairs Medical Center  
and Brown University

A deletion polymorphism in the serotonin transporter-linked polymorphic region (5-HTTLPR) has been associated with vulnerability to affective disorders, yet the mechanism by which this gene confers vulnerability remains unclear. Two studies examined associations between the 5-HTTLPR polymorphism and attentional bias for emotional stimuli among nondepressed adults. Biased attention, attention engagement, and difficulty with attention disengagement were assessed with a spatial cuing task using emotional stimuli. Results from Study 1 ( $N = 38$ ) indicated that short 5-HTTLPR allele carriers experienced greater difficulty disengaging their attention from sad and happy stimuli compared with long allele homozygotes. Study 2 participants ( $N = 144$ ) were genotyped for the 5-HTTLPR polymorphism, including single nucleotide polymorphism rs25531 in the long allele of the 5-HTTLPR. Consistent with Study 1, individuals homozygous for the low-expressing 5-HTTLPR alleles (i.e., S and  $L_G$ ) experienced greater difficulty disengaging attention from sad, happy, and fear stimuli than high-expressing 5-HTTLPR homozygotes. Because this association exists in healthy adults, it may represent a susceptibility factor for affective disorders that becomes problematic during stressful life experiences.

*Keywords:* genetic association, depression vulnerability, serotonin transporter, information processing

Individuals who inherit the low-expressing variant of the serotonin transporter-linked polymorphic region (5-HTTLPR) of the serotonin transporter gene (SLC6A4) and experience significant life stress appear to be at greater risk for depression than people who inherit two copies of the high-expressing 5-HTTLPR allele (e.g., Caspi et al., 2003; Kendler, Kuhn, Vittum, Prescott, & Riley, 2005). The landmark studies of Caspi et al. (2003) and Kendler et al. (2005) have stimulated research aimed at understanding why the low-expressing 5-HTTLPR variant putatively increases sensitivity to life stress and, in turn, increases vulnerability to depression. One possibility is that the 5-HTTLPR polymorphism biases the processing of emotional information. Therefore, we sought to examine whether healthy adult carriers of the low-expressing 5-HTTLPR allele display more pronounced attentional biases for emotional stimuli than individuals homozygous for the high-expressing 5-HTTLPR allele.

The serotonin transporter (5-HTT) regulates the reuptake of serotonin to the presynaptic neuron for recycling or degradation

after serotonin has been released. It thus plays a critical role in determining the duration and intensity of serotonin communication with postsynaptic receptors and targets, such as those in limbic regions involved in the regulation of emotional information (for a review, see Hariri & Holmes, 2006). Of importance is that the efficiency with which the 5-HTT returns serotonin to the presynaptic neuron appears to be influenced by the 5-HTTLPR polymorphism. A common deletion polymorphism in the promoter region of the 5-HTT gene results in two variants: a short (S) allele and a long (L) allele. The presence of one or two short alleles, rather than two copies of the long allele, is associated with reduced transcriptional efficiency that putatively results in significant decreases (approximately 50%) in serotonin reuptake (Heinz et al., 2000; Lesch et al., 1996). Short 5-HTTLPR allele carriers should thus have increased levels of extracellular serotonin and increased serotonin signaling compared with long allele homozygotes.

This difference in how serotonin is regulated appears to impact a cortical-limbic circuit that is critical for regulating emotional information. For instance, among healthy participants, Pezawas et al. (2005) found that short 5-HTTLPR allele carriers had significant reduction in gray matter volume in the perigenual anterior cingulate (pACC) and the rostral anterior cingulate—regions of the prefrontal cortex previously implicated in affect regulation (Drevets, 2000)—compared with long allele homozygotes. Furthermore, using functional magnetic resonance imaging (fMRI) analyses to assess relative activation of the pACC and amygdala in response to negative stimuli (e.g., angry and scared facial expressions), short 5-HTTLPR allele carriers had less functional coupling between the pACC and the amygdala. The “uncoupling” of this emotion circuit may explain why short-allele carriers have greater amygdala responses to emotional stimuli (Bertolino et al., 2005; Hariri et al., 2005, 2002).

---

Christopher G. Beevers, Tony T. Wells, and Alissa J. Ellis, Department of Psychology, University of Texas at Austin; John E. McGeary, Providence Veterans Affairs Medical Center and Center for Alcohol and Addiction Studies, Brown University.

Preparation of this article was facilitated by National Institute of Mental Health Grant R01MH076897 awarded to Christopher Beevers and National Center for Research Resources and the Department of Veteran Affairs shared equipment grants 1S10RR023457 awarded to John McGeary. We thank the research assistants from the Mood Disorders Laboratory at the University of Texas for their assistance with data collection.

Correspondence concerning this article should be addressed to Christopher G. Beevers, Department of Psychology, University of Texas at Austin, 1 University Station, A8000, Austin, TX 78712. E-mail: beevers@psy.utexas.edu

Other research has similarly found that reduced functional coupling between the prefrontal cortex and limbic system is associated with biased processing of emotional information at a neural level. Heinz et al. (2005) reported that a region of the prefrontal cortex more dorsal and rostral to the pACC was overactivated in short 5-HTTLPR allele carriers compared with long 5-HTTLPR homozygotes when presented with emotional images. Activation of the ventromedial prefrontal cortex region was positively correlated with amygdala activation, suggesting a compensatory effort to regulate exaggerated amygdala responses of the short allele carriers. This heightened amygdala response in short 5-HTTLPR allele carriers likely applies to a variety of emotional stimuli, as the amygdala also responds to positive stimuli (Cunningham, Van Bavel, & Johnsen, 2008), as well as to novel, salient, and ambiguous stimuli (Whalen, 2007).

Difficulty regulating emotional information at a neural level may have important implications for the cognitive processing of emotional information. The ability to allocate attention to emotional cues in the environment is a crucial element of adaptive self-regulation (Posner & Rothbart, 2000). Although it is adaptive for salient stimuli to capture attention, successful behavioral regulation requires some flexibility and control over attention. This may include strategic filtering of certain stimuli, timely disengagement from stimuli, and being appropriately vigilant for meaningful emotional cues. In line with this conclusion, Hasler, Drevets, Manji, and Charney (2004) argued that biased processing of emotional stimuli is a highly plausible and important putative intermediate phenotype for major depression disorder (MDD). Furthermore, they specifically identified the 5-HTTLPR as a highly promising candidate gene that may be associated with biased processing of emotional information (see Figure 1, p. 1766).

Despite this promising neuroimaging research, there have been few studies in which associations between the 5-HTTLPR polymorphism and behavioral assessments of biased processing of emotional information have been examined. Roiser, Cook, Cooper, Rubinsztein, and Sahakian (2005) reported that among users of ecstasy, a drug that causes long-term reductions in synaptic 5-HT release, short 5-HTTLPR allele carriers showed abnormal emotional processing (e.g., more likely to make errors of commission) than long 5-HTTLPR allele homozygotes on an affective go/no-go test. Similarly, Hayden et al. (2008) reported that children who were homozygous for the short 5-HTTLPR allele showed greater self-referent encoding of negative stimuli than the other 5-HTTLPR allele groups. Consistent with these analyses, work by Fallgatter and colleagues, using event-related potentials, indicated that short 5-HTTLPR allele carriers experience greater brain electrical activity during evaluation of an erroneous behavioral response than age- and gender-matched controls homozygous for the long 5-HTTLPR allele (Fallgatter et al., 2004).

We built on this research in the present study and provided an additional test of whether the 5-HTTLPR polymorphism is associated with biased processing of emotional stimuli. In previous work, using a standard dot probe task, Beevers, Gibb, McGeary, and Miller (2007) reported that short 5-HTTLPR allele carriers displayed biased attention for anxious-relevant words when paired with neutral words compared with long 5-HTTLPR allele homozygotes. Although those findings were intriguing, that study had a number of limitations. Participants were a heterogeneous mix of psychiatric inpatients experiencing a variety of emotional states

and pharmacological treatments, which may have influenced the serotonin system and emotion processing. Furthermore, there are concerns that the dot probe task, especially when used with longer stimulus durations, may not directly assess attentional engagement but might instead assess difficulty with the disengagement of attention. As a result, researchers have suggested using tasks in which attention engagement and disengagement are measured separately (e.g., Koster, De Raedt, Goeleven, Franck, & Crombez, 2005; Posner & Petersen, 1990).

We therefore used a spatial cuing task that allowed us to assess general attentional bias, attentional engagement, and difficulty with attentional disengagement from emotional stimuli (Koster et al., 2005). For this version of the task, a cue is presented to the left or right side of visual field. Cues are faces expressing an emotion (e.g., happiness, sadness) or no emotion (e.g., neutral). After the offset of the cue, a target appears either in the same location of the cue (e.g., valid cue trial) or in the opposite side of visual field (e.g., invalid cue trial). The dependent measure is latency to identify the location of the cue. Typically, reaction times are shorter for valid cue trials than for invalid cue trials, although this pattern can reverse when cues are presented for longer durations (i.e., inhibition of return effect). The difference in mean reaction time for invalid and valid trials is referred to as the *cue validity effect*. A positive cue validity effect is interpreted as maintained attention to the cue.

With the emotional modification of the spatial cuing task, it is possible to determine whether the cue validity effect is influenced by cue valence (e.g., happy, sad, neutral facial expressions). Three different but related outcomes can be examined: (a) general measure of biased attention, (b) attentional engagement, and (c) difficulty with attentional disengagement. Each of these outcomes compares reaction time to identify targets following emotional cues relative to neutral cues. For instance, if the cue validity effect was stronger for emotional cues than neutral cues, we would observe a positive biased attention score for emotional stimuli. This would indicate that attention was maintained by emotional cues to a greater extent than by neutral cues (Mogg, Holmes, Garner, & Bradley, 2008). Similarly, faster latencies following emotional valid cues than neutral valid cues suggests that emotional stimuli facilitated attentional engagement, whereas longer latencies following invalid emotional cues compared with invalid neutral cues suggests difficulty disengaging attention from emotional cues. Together, these outcomes provide a general assessment of maintained attention and the specific attentional processes related to maintained attention (Koster et al., 2005).

Using the spatial cuing task, we conducted two studies designed to address the limitations of our previous study and provide a more precise test of the hypothesis that the 5-HTTLPR polymorphism is associated with attentional biases for emotional stimuli. Specifically, in Study 1, we recruited a sample of healthy adults who did not meet criteria for a current psychiatric disorder or a past history of a mood disorder and who were not currently taking any psychiatric medications. These individuals were genotyped for the 5-HTTLPR polymorphism and then completed a spatial cuing task in which sad, happy, and neutral facial expressions were used as cues. For Study 2, we recruited a large sample of unmedicated, nondepressed undergraduate students. We administered the same spatial cuing task as in Study 1 but also included fear facial expressions as cues.

We expected to observe an interaction between cue validity, stimulus valence, and 5-HTTLPR allele group when predicting response time latencies on the spatial cuing task. We hypothesized that carriers of the low-expressing 5-HTTLPR allele would display biased attention for emotional stimuli in general; however, given the lack of specificity with which we measured biased attention in our previous work, it was unclear whether we should expect enhanced attentional engagement or increased difficulty with attentional disengagement among the low-expressing 5-HTTLPR allele carriers. Therefore, we did not make specific hypotheses regarding each component of biased attention.

## Study 1

### Method

#### Participants

Participants were 38 adults recruited from the Austin, Texas, community (see Table 1 for demographic information). On average, the sample was in their early 30s, Caucasian, female, and educated. Participants were recruited using flyers posted in the community and with ads posted on electronic Web sites. Participants received \$15 an hour for their participation. The average laboratory session was 2 hr. Interviewers used structured clinical interviews to determine presence and history of psychopathology. Inclusion criteria included normal or corrected-to-normal vision, fluency in the English language, and age between 22 and 60 years. Exclusion criteria included the presence of any current Axis I disorder. Furthermore, participants who had a history of MDD or dysthymia, or were currently prescribed psychiatric medication were also excluded.

#### Assessments

*Mini International Neuropsychiatric Interview (MINI).* The electronic version of the MINI was used as part of a screening interview to determine whether participants met criteria for study entry. The MINI is a short, structured screening interview that was developed for the *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition (*DSM-IV*; American Psychiatric Association, 1994) and the International Classification of Diseases, 10th

edition (ICD-10) psychiatric disorders (Kendler et al., 2005; Sheehan et al., 1998). The MINI assesses diagnostic criteria for the following disorders: MDD, dysthymia, panic disorder, agoraphobia, social phobia, obsessive-compulsive disorder, posttraumatic stress disorder, psychotic disorder, anorexia nervosa, bulimia nervosa, generalized anxiety disorder, and antisocial personality disorder. It also assesses for suicidal ideation and behavior, mania, and hypomania. The MINI has been validated against the Structured Clinical Interview for *DSM-IV* (SCID; First, Spitzer, Gibbon, & Williams, 2002) diagnoses and against the Composite International Diagnostic Interview for ICD-10 (Lecrubier et al., 1997; Sheehan et al., 1998).

Interviewers were undergraduate research assistants who received at least 10 hr of training, wherein they learned interview skills, reviewed diagnostic criteria, and role-played interviews. Because this was a screening interview, brevity was important. Interviewers could terminate the interview as soon as the participant did not meet study criteria. Therefore, the entire MINI was typically completed only for participants who met criteria for study entry. The average length of MINI screening interview was approximately 15 min.

*SCID.* To confirm key inclusion/exclusion criteria from the screening interview, the Mood Disorders and the Anxiety Disorders Modules of the patient version of the SCID (First et al., 2002) were administered during an in-person interview at the time of study participation. Three assessors with at least a bachelor's degree in psychology conducted all interviews. Assessors participated in 15 hr of training, wherein they learned interview skills, reviewed diagnostic criteria for relevant *DSM-IV* diagnoses (American Psychiatric Association, 1994), observed mock interviews, and role-played interviews. Twenty percent of all interviews were rated by an independent assessor. Agreement between study interviewer and independent assessor diagnosis of MDD was excellent ( $k = 1.0$ ).

*Beck Depression Inventory-II (BDI-II; Beck, Brown, & Steer, 1996).* The BDI-II is a widely used self-report questionnaire that assesses depression severity. The BDI-II consists of 21 items and measures the presence and severity of cognitive, motivational, affective, and somatic symptoms of depression. Past reports have indicated test-retest reliability is adequate (Beck, Steer, & Garbin,

Table 1  
*Study 1 Demographics as a Function of 5-HTTLPR Allele Status*

Demographic	LL ( $n = 14$ )	SL ( $n = 17$ )	SS ( $n = 7$ )
Age (years)	28.79 (7.77)	30.07 (6.65)	32.86 (6.59)
Gender (M/F)	21%/79%	18%/82%	43%/57%
Hispanic (yes/no)	7%/93%	11%/89%	28%/72%
Race (Caucasian/"other")	93%/7%	77%/23%	72%/28%
Married (yes/no)	36%/64%	24%/76%	28%/72%
College degree (yes/no)	64%/36%	59%/41%	85%/15%
Income (mean)	\$38,500	\$50,400	\$39,142
Depressive symptoms (BDI-II)	2.64 (4.19)	6.07 (8.82)	4.86 (6.07)

*Note.* Means and standard deviations appear for Age and Depressive symptoms. 5-HTTLPR = serotonin transporter-linked polymorphic region; LL = long/long; SL = short/long; SS = short/short; M = male; F = female; BDI-II = Beck Depression Inventory-II.

1988). The BDI-II has been found to be valid among psychiatric inpatient and outpatient samples (Beck et al., 1988). Internal consistency reliability in the present study was good ( $\alpha = .81$ ).

**Genotyping.** Genomic DNA was isolated from buccal cells using a modification of published methods (Freeman et al., 1997; Lench, Stanier, & Williamson, 1988; Meulenbelt, Droog, Trommelen, Boomsma, & Slagboom, 1995; Spitz et al., 1996). The cheeks and gums are rubbed for 20 s with three sterile, cotton-tipped wooden swabs. The swabs are placed in a 50-ml capped polypropylene tube containing lysis buffer (500  $\mu$ l of 1 M Tris-HCl; 200 mM disodium ethylene diaminetetracetic acid [EDTA], pH 8.0; 500  $\mu$ l of 10% sodium dodecyl sulfate; and 100  $\mu$ l of 5 M sodium chloride). The participants then rinse out the mouth vigorously with 10 ml of distilled water for 20 s, and this was added to the 50-ml tube. The tubes were stored at 4 °C until the DNA was extracted.

To extract the DNA, proteinase K (0.2mg/ml) was added to the tubes, and the tubes were incubated at 65 °C for 60 min. The swabs were removed and residual lysis buffer was extracted by centrifugation (using a 3-ml syringe barrel and sterile 15-ml tube) for 5 min at 1,000  $\times$  g. The residual fluid was added back to the original sample. An equal volume of isopropyl alcohol was then added to each tube, the contents were mixed, and the DNA was collected by centrifugation at 3,500  $\times$  g for 10 min. The DNA pellet was rinsed once with 1 ml of 50% isopropyl alcohol and allowed to air dry. The pellet was resuspended in 1 ml of 10  $\mu$ M TRIS-HCl, 10 mM EDTA buffer (pH 8.0) and placed in a 1.8-ml cryovial.

The 5-HTT gene (SLC6A4), which maps to 17q11.1–17q12, contains a 44-bp deletion in the 5-in. (13-cm) regulatory region of the gene (Heils et al., 1996). The variable number of tandem repeats in the promoter appears to be associated with variations in transcriptional activity: the long (L) variant has approximately three times the basal activity of the shorter (S) promoter with the deletion (Lesch et al., 1996). The assay is a modification of the method of Lesch and colleagues (Lesch et al., 1996).

**Amplification of 5-HTTLPR target sequences by polymerase chain reaction.** Polymerase chain reactions (PCRs) contain 0.5  $\mu$ l of genomic DNA (20 ng), 10% DMSO (Hybra-Max® grade; Sigma-Aldrich, St. Louis, MO), 1.8 mM MgCl<sub>2</sub>, 180  $\mu$ M deoxynucleotides, with 7'-deaza-2'-deoxyGTP (Roche Applied Science, Indianapolis, IN) substituted for one half of the deoxyguanosine triphosphate, forward and reverse primers (5-HTT, 480 nM; obtained from IDT, Coralville, IA) and 1 unit (U) of AmpliTaq Gold® polymerase (Applied Biosystems, Foster City, CA), in a total volume of 20  $\mu$ l. Amplification is performed using touchdown PCR (Don, Cox, Wainwright, Baker, & Mattick, 1991). A 95 °C incubation for 10 min is followed by two cycles of 95 °C for 30 s, 65 °C for 30 s, and 72 °C for 60 s. The annealing temperature is decreased every two cycles from 65 °C to 57 °C in 2°-C increments (10 cycles total), and a final 30 cycles of 95 °C for 30 s, 55 °C for 30 s, and 72 °C for 60 s, and a final 30-min incubation at 72 °C. After amplification, 1  $\mu$ l of PCR product is combined with 2  $\mu$ l of loading buffer containing size standard (Genescan 2500 TAMRA®; Applied Biosystems), and 0.8  $\mu$ l is loaded into each of 48 lanes of a 12-cm gel. PCR products are electrophoresed through a 4.25% polyacrylamide gel under denaturing conditions (6 M urea) with an Applied Biosystems ABI Prism 377 DNA sequencer using protocols supplied by the company.

The primer sequences are the following: forward, 5'-GGCGTTGCCGCTCTGAATGC-3' (fluorescently labeled), and reverse, 5'-GAGGGACTGAGCTGGACAACCAC-3'. These primer sequences yield products of 484 or 528 bp. Two investigators scored allele sizes independently, and any inconsistencies were reviewed and rerun (less than 20% of sample). An additional 10% of the sample was randomly selected and rerun to ensure genotyping accuracy. There was 100% agreement for allele groupings between the initial and retested samples. The 5-HTTLPR allele frequency distribution was SS:  $n = 7$  (18.4%), SL:  $n = 17$  (44.7%), LL:  $n = 14$  (36.8%). Genotype distribution was in Hardy-Weinberg equilibrium,  $\chi^2(1, N = 38) = 0.207, p = .649$ , and is consistent with previous reports (Lesch et al., 1996).

**Spatial cuing task.** The spatial cuing task was developed by Posner (1980) and modified by others to incorporate emotional cues (e.g., Koster et al., 2005). Each trial sequence (shown in Figure 1) began by presenting a fixation cross in the center of the screen for 500 ms. Then, a face cue was presented on either the left or the right side of the visual field for 1,500 ms. After cue offset, a probe (either \* or \*\*) immediately appeared on the left or right side of visual field and remained on the screen until the participant responded by pressing an appropriate response box button. The participant's task was to identify probe type as quickly and accurately as possible. Participants pressed a corresponding button on a response box to indicate the type of probe that appeared. After the participant responded, the screen was blank for 500 ms before the next trial began. Seventy-five percent of probes appeared on the same side of visual field as the cue (a valid trial). Twenty-five percent of the probes appeared on the opposite side of visual field as the cue (an invalid trial). Both valid and invalid trials had a 50% chance of having either the single- or double-asterisk probe.

Cue stimuli were images of faces taken from the Pictures of Facial Affect (Ekman & Friesen, 1976) photo set. Human faces were selected because facial expressions receive special processing priority (Farah, Wilson, Drain, & Tanaka, 1998), and because human faces have been used extensively in behavioral and imaging studies, and are arguably more ecologically valid than written words. Twelve faces were selected from each of the following categories: happy, sad, and neutral. All stimuli were presented on a black background on a 17-in. (43-cm) color monitor. Stimuli were approximately 10.5  $\times$  17 cm when presented on the screen. Participants completed 10 practice trials using neutral faces as




Figure 1. Trial sequence for valid and invalid trials. Fixation cross and probe are not to scale.

cues. Anyone failing to respond accurately to at least 8 of the 10 trials repeated the practice trials until they had achieved 80% accuracy. Participants then completed a total of 72 trials. They viewed each of the 36 stimuli twice. Order of stimulus presentation was randomized for each participant, with the stipulation that each of the 36 stimuli were viewed once before stimuli were repeated.

Of interest were the three primary outcomes from the spatial cuing task: general attentional bias, attentional engagement, and difficulty with attentional disengagement. As suggested by Mogg et al. (2008), a general measure of attentional bias can be derived from the spatial cuing task using the following formula:

Attentional bias score

$$= (\text{RT invalid emotion cue trials} - \text{RT valid emotion cue trials}) \\ - (\text{RT invalid neutral cue trials} - \text{RT valid neutral cue trials}). \quad (1)$$

This attentional bias score does not attempt to differentiate attentional engagement versus disengagement. Positive values reflect an attentional bias for emotional cues relative to neutral cues. Negative values reflect an attentional bias for neutral cues relative to emotional cues.

Attentional engagement and difficulty with attentional disengagement can be calculated for each of the emotion cue categories with the following formulae:

Attentional engagement =

$$\text{RT valid neutral cue} - \text{RT valid emotional cue}. \quad (2)$$

Difficulty with attentional disengagement =

$$\text{RT invalid emotional cue} - \text{RT invalid neutral cue}. \quad (3)$$

Positive attentional engagement scores indicate that participants identify the target more quickly when it follows an emotion cue from that category than when it follows a neutral cue. Negative scores indicate the reverse. Positive attentional disengagement scores indicate greater difficulty disengaging attention from an emotional cue than from a neutral cue. That is, positive scores indicate a slower shift of attention away from an emotional cue than from a neutral cue. Negative scores indicate less difficulty disengaging attention from emotional cues than from a neutral cue. Mean reaction time scores are used to compute each of these outcomes.

### Procedure

Participants contacted the Mood Disorders Laboratory at the University of Texas at Austin expressing an interest in study participation. The MINI screening interview was then conducted by a trained interviewer. If the participant passed the screening assessment, they were scheduled for a laboratory appointment. Upon arrival to the laboratory, participants were oriented to the laboratory, provided informed consent, and then completed a demographics form. They next completed the SCID interview to confirm the absence of mood and anxiety disorders. Participants then completed several self-report questionnaires and provided buccal cells via a cheek swab/mouthwash procedure for genotyp-

ing. Next, participants completed the spatial cuing task (and other tasks not included in this report). Upon completion of study procedures, participants were debriefed and paid \$15 an hour (up to a maximum of \$50) for their participation. The Internal Review Board at the University of Texas at Austin approved all study procedures.

## Results

### Sample Characteristics

Descriptive statistics for the sample are presented in Table 1 stratified by 5-HTTLPR allele status. There were no significant differences as a function of allele grouping for age,  $F(2, 37) = 0.58, p = .57$ ; gender,  $\chi^2(2, N = 38) = 4.26, p = .12$ ; ethnicity,  $\chi^2(2, N = 38) = 1.93, p = .38$ ; race,  $\chi^2(2, N = 38) = 1.93, p = .38$ ; marital status,  $\chi^2(2, N = 38) = 0.55, p = .76$ ; education,  $\chi^2(2, N = 38) = 1.61, p = .44$ ; income,  $F(2, 37) = 0.58, p = .57$ ; and depressive symptoms,  $F(2, 37) = 0.65, p = .53$ .

### Data Reduction

We deleted trials with incorrect responses (0.4% of all trials) and did not use them for analyses. Furthermore, we deleted reaction times that were at least three standard deviations beyond the individual reaction time mean (1.4%). Together, these procedures resulted in the exclusion of less than 1.9% of the data.

*Main results.* We conducted a 2 (cue validity: valid, invalid)  $\times$  3 (stimulus valence: sad, happy, neutral)  $\times$  3 (allele status: SS, SL, LL) mixed-plot analysis of variance (ANOVA) to examine whether the 5-HTTLPR polymorphism was associated with latency to identify target (see Table 2 for descriptive statistics). Results indicated a significant cue validity main effect,  $F(1, 35) = 26.39, p < .001, \eta^2 = .43$ , and a Stimuli Valence  $\times$  Allele Status interaction that approached significance,  $F(4, 70) = 2.15, p = .08, \eta^2 = .11$ . The Cue Validity  $\times$  Allele Status interaction,  $F(2, 35) = 0.95, p = .40, \eta^2 = .05$ ; the stimulus valence main effect,  $F(2, 70) = 1.70, p = .19, \eta^2 = .05$ ; the Cue Validity  $\times$  Stimulus Valence,  $F(2, 70) = 0.02, p = .98, \eta^2 = .00$ ; and the Cue Validity  $\times$  Stimulus Valence  $\times$  Allele Status interaction,  $F(4, 70) = 1.11, p = .36, \eta^2 = .06$ , did not reach statistical significance. However, given the preliminary nature of Study 1, and that conducting a replication study (Study 2) with a larger sample would help to tease apart Type I errors from true effects, we examined whether 5-HTTLPR allele status was associated with biased attention, attentional engagement, and attentional disengagement.

*Attentional bias.* We conducted a 2 (stimulus valence: sad, happy)  $\times$  2 (allele status: SS, SL, LL) mixed-plot ANOVA to examine whether 5-HTTLPR allele status predicted attentional bias scores. Results indicated a nonsignificant main effect for valence,  $F(1, 35) = 0.002, p = .96, \eta^2 = .00$ , and a nonsignificant Valence  $\times$  Allele Status interaction,  $F(2, 35) = 0.26, p = .78, \eta^2 = .01$ . However, the main effect for allele status approached significance,  $F(2, 35) = 2.50, p = .09, \eta^2 = .12$ . Pairwise least significant difference (LSD) comparisons indicated significant differences between the LL and SS groups ( $p = .05$ ), marginal difference between the LL and SL groups ( $p = .09$ ), and no differences between the SL and SS groups ( $p = .56$ ). Mean

Table 2

Means (and Standard Deviations) for Latencies to Identify Target (in milliseconds) Presented by Cue Validity, Stimulus Valence, and 5-HTTLPR Allele Status for Study 1

Stimulus valence	5-HTTLPR allele group					
	LL		SL		SS	
	Valid cue	Invalid cue	Valid cue	Invalid cue	Valid cue	Invalid cues
Happy	645 (97)	662 (110)	702 (205)	766 (231)	616 (118)	667 (146)
Sad	634 (117)	661 (92)	696 (199)	750 (212)	619 (111)	672 (135)
Neutral	633 (94)	684 (120)	679 (197)	728 (201)	617 (126)	643 (127)

Note. 5-HTTLPR = serotonin transporter-linked polymorphic region; LL = long/long; SL = short/long; SS = short/short.

attentional bias scores (and standard errors) collapsing across stimulus valence were LL:  $-29.23$  ( $19.59$ ), SL:  $9.97$  ( $12.20$ ), SS:  $26.28$  ( $22.28$ ). Figure 2A presents attentional bias scores for each stimulus valence.

**Attentional engagement and disengagement.** We used a 2 (bias: engagement, disengagement)  $\times$  2 (stimulus valence: sad, happy)  $\times$  3 (allele status: S, SL, LL) mixed-plot ANOVA to examine whether the 5-HTTLPR polymorphism was associated with attentional engagement and disengagement for emotional stimuli. Results indicated a significant interaction between bias and allele status,  $F(2, 35) = 3.63$ ,  $p < .05$ ,  $\eta^2 = .17$ . The bias main effect,  $F(1, 35) = 2.71$ ,  $p = .11$ ,  $\eta^2 = .07$ ; stimulus valence main effect,  $F(1, 35) = 0.02$ ,  $p = .96$ ,  $\eta^2 = .00$ ; Stimulus Valence  $\times$

Allele Status interaction,  $F(2, 35) = 0.26$ ,  $p = .78$ ,  $\eta^2 = .01$ ; Bias  $\times$  Stimulus Valence interaction,  $F(1, 35) = 0.53$ ,  $p = .47$ ,  $\eta^2 = .02$ ; and the three-way interaction,  $F(2, 35) = 0.66$ ,  $p = .66$ ,  $\eta^2 = .02$ , did not reach statistical significance.

We followed up the significant interaction by examining attentional engagement and disengagement scores across the genetic groups. As there were no significant effects involving stimulus valence, we collapsed across valence for these follow-up analyses. For attentional engagement, results of a univariate ANOVA (allelic status: S, SL, LL) revealed no significant differences between 5-HTTLPR allele groups,  $F(1, 35) = 1.17$ ,  $p = .13$ ,  $\eta^2 = .06$  (see Figure 2B). For attentional disengagement, there was a significant main effect for allele status,  $F(2, 35) = 3.74$ ,  $p < .05$ ,  $\eta^2 = .18$ .




Figure 2. A: Mean attentional bias, B: attentional engagement, and C: attentional disengagement, with standard errors presented as a function of stimulus valence and serotonin transporter-linked polymorphic region (5-HTTLPR) allele status for Study 1. SS = short/short; SL = short/long; LL = long/long.

Pairwise LSD comparisons indicated that the LL group disengaged from emotional stimuli significantly faster than the SL group ( $p = .02$ ) and marginally faster than the SS group ( $p = .06$ ). There was no significant difference in disengagement between the SS and SL groups ( $p = .86$ ; see Figure 2C).

Finally, we also examined whether attentional engagement and disengagement scores (collapsing across valence) were significantly different from zero for each of the genetic groups. For engagement scores, only the SL group was significantly different from zero,  $t(16) = -2.27, p < .05$ . For disengagement scores, the SL group was also significantly greater than zero,  $t(16) = 2.13, p < .05$ , whereas the SS,  $t(6) = 1.47, p = .17$ , and LL,  $t(13) = -1.47, p = .16$ , groups did not significantly differ from zero.

### Discussion

Findings from Study 1 provide tentative evidence that short 5-HTTLPR allele carriers exhibit biased attention for emotional stimuli compared with individuals homozygous for the long 5-HTTLPR allele. Furthermore, the 5-HTTLPR polymorphism was associated with difficulty disengaging attention from happy and sad facial expressions relative to neutral faces but was not associated with attentional engagement. This is among the first evidence that the 5-HTTLPR polymorphism influences attentional biases for emotional stimuli among healthy controls.

Nevertheless, we thought it was important to replicate and extend these initial findings for several reasons. First, initial behavioral genetics findings are often not replicated in subsequent studies, particularly when the initial studies involve small samples (Ioannidis, Ntzani, Trikalinos, & Contopoulos-Ioannidis, 2001). This concern is particularly relevant for the present study, as the predicted three-way interaction between cue validity, stimulus valence, and 5-HTTLPR allele status was not statistically significant. Thus, analyses of linkages between the 5-HTTLPR polymorphism and biased attention are likely associated with a higher Type I error rate than is typically tolerated in a single study. We therefore conducted a second test of the hypothesized three-way interaction in a larger sample of participants in Study 2.

Second, it has recently been discovered that the long 5-HTTLPR allele may have two variants (i.e.,  $L_A$  and  $L_G$ ). In the first of two extra 20- to 23-bp repeats in the L allele, a common single nucleotide polymorphism occurs at the sixth nucleotide (adenine to guanine; A to G). More important, the  $L_G$  variant and the S allele appear to be very similar in terms of transcriptional activity; therefore, only the  $L_A$  variant is high expressing with regard to

transcriptional activity (Hu et al., 2005). In Study 2, we used this more recent approach to genotype the 5-HTTLPR and more precisely examine the association of the 5-HTTLPR genotype with biased attention.

Third, we added a third emotional stimulus category to the spatial cuing task, faces depicting fear. We included this stimulus category to determine whether carriers of the low-expressing 5-HTTLPR allele display greater biased attention and experience greater difficulty disengaging their attention from negative stimuli other than sad stimuli. We therefore conducted a second study with the modifications described above to replicate and extend our initial findings.

### Study 2

For Study 2, we recruited a convenient sample of nondepressed undergraduate students who provided buccal cells for genetic analyses and then completed the spatial cuing task. On the basis of the findings from Study 1, we expected carriers of the low-expressing 5-HTTLPR allele to exhibit greater biased attention for emotional stimuli than individuals homozygous for the high-expressing 5-HTTLPR allele. Furthermore, we also expected 5-HTTLPR allele group differences for difficulty disengaging attentional, but none were expected for attentional engagement.

### Method

#### Participants

Participants were 144 nondepressed individuals recruited from introductory psychology classes at the University of Texas at Austin. All participants had BDI-II scores of 9 or fewer and were unmedicated at the time of testing (see Table 3 for descriptive statistics). Participants received course credit for their participation. This study took approximately 30 min to complete.

#### Genotyping

DNA samples were genotyped for the 5-HTTLPR deletion polymorphism using methods described in Study 1. The 5-HTTLPR allele frequency distribution was SS:  $n = 28$  (19.4%), SL:  $n = 73$  (50.6%), LL:  $n = 43$  (29.9%). Genotype distribution was in Hardy-Weinberg equilibrium,  $\chi^2(1, N = 144) = 0.09, p = .76$ . To distinguish between the S,  $L_A$ , and  $L_G$  fragments, the PCR fragment was digested with MspI according to the methods found

Table 3  
Study 2 Demographics as a Function of 5-HTTLPR Allele Status

Demographic	L'L' ( $n = 31$ )	S'L' ( $n = 84$ )	S'S' ( $n = 29$ )
Age (years)	19.31 (1.71)	19.02 (1.11)	18.94 (1.10)
Gender (M/F)	57%/43%	40%/60%	47%/53%
Hispanic (yes/no)	15%/84%	17%/83%	26%/74%
Race (Caucasian/"other")	82%/18%	70%/30%	60%/40%
Marital status (married: yes/no)	6%/94%	4%/96%	3%/97%
Depressive symptoms (BDI-II)	3.77 (2.21)	2.88 (2.66)	1.78 (2.44)

Note. S'S' includes the SS,  $L_G L_G$ , and  $SL_G$  allele groups, S'L' includes the  $SL_A$ ,  $L_G L_A$  allele groups, and the L'L' includes the  $L_A L_A$  allele groups. 5-HTTLPR = serotonin transporter-linked polymorphic region; BDI-II = Beck Depression Inventory-II. Means and standard deviations appear for Age and Depressive symptoms.

in Wigg et al. (2006). The resulting polymorphic fragments were separated using an ABI 377 DNA sequencer (S: 297, 127, and 62 bp; L<sub>A</sub>: 340, 127, and 62 bp; L<sub>G</sub>: 174, 166, 127, and 62 bp). Using this approach, allele frequencies were S:  $n = 129$  (44.7%), L<sub>A</sub>:  $n = 146$  (50.6%), L<sub>G</sub>:  $n = 13$  (4.5%). Genotype distribution for the A/G single nucleotide polymorphism was in Hardy-Weinberg equilibrium,  $\chi^2(1, N = 144) = 0.32, p = .57$ .

Consistent with previous research (Hu et al., 2005; Zalsman et al., 2006), the low-expressing S and L<sub>G</sub> alleles were designated S', and the higher expressing L<sub>A</sub> allele was designated L'. Three groups were formed: (a) S'S' (i.e., SS:  $n = 28$  [19.4%], SL<sub>G</sub>:  $n = 1$  [0.7%], L<sub>G</sub>L<sub>G</sub>:  $n = 0$  [0%]), (b) S'L' (i.e., SL<sub>A</sub>:  $n = 72$  [50.0%], L<sub>G</sub>L<sub>A</sub>:  $n = 12$  [8.3%]), and (c) L'L' (i.e., L<sub>A</sub>L<sub>A</sub>:  $n = 31$  [21.5%]).

### Procedure

Undergraduate students who scored less than 4 on the short-form of the BDI (BDI-SF) during mass pretesting were invited to participate in this study. Upon arrival to the laboratory, depression severity was reassessed using the BDI-II. Participants with BDI-II scores greater than 9 or anyone currently taking psychiatric medications were excluded from this study. In contrast to Study 1, participants did not complete a SCID interview; however, all participants were experiencing low levels of depressive symptoms. Participants who qualified were then oriented to the laboratory, provided informed consent, and completed a demographics form. Participants completed several self-report questionnaires and provided buccal cells via a cheek swab/mouthwash for genotyping. Next, participants completed the spatial cuing task. The Internal Review Board at the University of Texas at Austin approved all study procedures.

**Spatial cuing task.** The task was the same as in Study 1, with two important exceptions. First, 12 fear stimuli were included in addition to the sad, happy, and neutral stimuli used in Study 1. These stimuli were also selected from the Ekman and Friesen (1976) collection. Furthermore, participants were instructed to identify the location of the probe (left or right side of visual field) rather than the probe type. This change was made because probe location instructions can lead to fewer participant errors and produce results that are highly similar to probe identification instructions (Mogg & Bradley, 1999). The present task thus required participants to complete 96 trials (12 stimuli  $\times$  4 emotion categories  $\times$  2 presentations). As before, 75% of the trials were validly

cued trials, and 25% were invalidly cued trials. The probe was equally likely to appear on the left or the right side of visual field following a valid and invalid trial.

## Results and Discussion

### Sample Characteristics

Descriptive statistics for the sample are presented in Table 3 stratified by 5-HTTLPR allele status. There were no significant differences as a function of allele grouping for age,  $F(2, 143) = 1.57, p = .21$ ; gender,  $\chi^2(2, N = 143) = 2.93, p = .23$ ; ethnicity,  $\chi^2(2, N = 143) = 2.43, p = .29$ ; marital status,  $\chi^2(2, N = 143) = 1.80, p = .41$ ; and depressive symptoms,  $F(2, 143) = 2.78, p = .11$ . There were significant allele frequency differences for race,  $\chi^2(2, N = 143) = 13.48, p < .01$ . Therefore, we entered race as a covariate in all subsequent analyses; however, race was not significantly associated with any of the attentional bias outcomes (see the *Main Results* section).

### Data Reduction

We deleted and did not use trials with incorrect responses (0.5% of all trials) for analyses. Furthermore, we deleted reaction times that were at least three standard deviations beyond the individual reaction time mean (0.9%). Together, these procedures resulted in the exclusion of less than 1.5% of the data.

### Main Results

We used a 2 (cue validity: valid, invalid)  $\times$  4 (stimulus valence: sad, happy, fear, neutral)  $\times$  3 (5-HTTLPR allele type: S'S', S'L', L'L') mixed-plot ANOVA to examine whether the 5-HTTLPR polymorphism was associated with latency to identify the target (see Table 4 for descriptive statistics). Results indicated a significant three-way interaction between stimulus valence, cue validity, and 5-HTTLPR allele status,  $F(6, 420) = 2.32, p = .03, \eta^2 = .03$ . The cue validity main effect,  $F(1, 140) = 0.12, p = .73, \eta^2 = .00$ ; Cue Validity  $\times$  Allele Type interaction,  $F(2, 141) = 1.07, p = .35, \eta^2 = .00$  stimulus valence main effect,  $F(3, 420) = 1.39, p = .24, \eta^2 = .01$ ; Stimulus Valence  $\times$  Allele Group interaction,  $F(6, 420) = 0.94, p = .46, \eta^2 = .01$ ; and the Stimulus Valence  $\times$  Cue Validity interaction,  $F(3, 420) = 0.75, p = .94, \eta^2 = .00$ , did

Table 4

*Means (and Standard Deviations) Latencies in Milliseconds to Identify Target Presented by Cue Validity, Stimulus Valence, and 5-HTTLPR Allele Status for Study 2*

Stimulus valence	5-HTTLPR allele group					
	L'L'		S'L'		S'S'	
	Valid cue	Invalid cue	Valid cue	Invalid cue	Valid cue	Invalid cue
Happy	419 (61)	406 (59)	420 (52)	419 (59)	431 (74)	447 (83)
Sad	416 (56)	412 (59)	420 (59)	423 (72)	437 (74)	446 (86)
Fear	418 (57)	413 (55)	423 (52)	420 (58)	441 (87)	442 (85)
Neutral	418 (59)	419 (51)	418 (54)	421 (66)	436 (85)	431 (80)


*Note.* 5-HTTLPR = serotonin transporter-linked polymorphic region; L'L' = the L<sub>A</sub>L<sub>A</sub> allele groups; S'L' = the SL<sub>A</sub>, L<sub>G</sub>L<sub>A</sub> allele groups; S'S' = the SS, L<sub>G</sub>L<sub>G</sub>, and SL<sub>G</sub> allele groups.

not reach statistical significance. Furthermore, the race main effect,  $F(1, 140) = 0.70, p = .98, \eta^2 = .00$ ; Race  $\times$  Stimulus Valence interaction,  $F(3, 420) = 1.21, p = .30, \eta^2 = .01$ ; Cue Validity  $\times$  Race interaction,  $F(1, 140) = 0.53, p = .47, \eta^2 = .00$ ; Stimulus Valence  $\times$  Cue Validity  $\times$  Race interaction,  $F(3, 420) = 0.32, p = .81, \eta^2 = .00$ , were all nonsignificant. We followed up the significant three-way interaction by examining attentional bias, attentional disengagement, and attentional disengagement, computed as described in Study 1.

**Attentional bias.** We conducted a 3 (stimulus valence: sad, happy, fear)  $\times$  2 (allele status: S'S', S'L', L'L') mixed-plot ANOVA to examine whether 5-HTTLPR allele status predicted attentional bias scores. Results indicated a nonsignificant main effect for stimulus valence,  $F(2, 282) = 0.60, p = .55, \eta^2 = .00$ , and a Stimulus Valence  $\times$  Allele Status interaction that approached significance,  $F(4, 282) = 2.02, p = .09, \eta^2 = .03$ . However, the main effect for allele status was significant,  $F(2, 141) = 3.01, p = .05, \eta^2 = .04$ . Pairwise LSD comparisons indicated significant group differences between the L'L' and S'S' groups ( $p = .04$ ), no difference between the L'L' and S'L' groups ( $p = .86$ ), and a significant group difference between the S'L' and S'S' groups ( $p = .02$ ). Mean attentional bias scores (and standard errors) collapsing across stimulus valence were L'L':  $-6.41 (6.84)$ , S'L':  $-5.01 (4.16)$ , SS:  $13.97 (7.08)$ . Thus, the S'S' group showed a significantly greater attentional bias than the L'L' and S'L' groups. Figure 3A presents attentional bias scores for each stimulus valence.

**Attentional engagement and disengagement.** We conducted a 2 (bias: engagement, disengagement)  $\times$  3 (stimulus valence: sad, happy, fear)  $\times$  3 (allele status: S'S', S'L', L'L') mixed-plot ANOVA to examine whether 5-HTTLPR polymorphism was associated with attentional engagement and disengagement for emotional stimuli. Results indicated a significant Stimulus Valence  $\times$  Allele Status interaction,  $F(4, 282) = 2.67, p = .03, \eta^2 = .04$ , and a significant Bias  $\times$  Allele Status interaction,  $F(2, 141) = 3.82, p < .05, \eta^2 = .05$ . The stimulus valence main effect,  $F(2, 282) = 0.45, p = .77, \eta^2 = .00$ ; bias main effect,  $F(1, 140) = 0.42, p = .51, \eta^2 = .00$ ; Stimulus Valence  $\times$  Bias interaction,  $F(2, 282) = 1.18, p = .31, \eta^2 = .01$ ; and the three-way interaction were not significant,  $F(4, 282) = 0.45, p = .77, \eta^2 = .01$ .

We followed up the significant Bias  $\times$  Allele Status interaction by examining reaction time for each attention bias type (collapsing across stimulus valence) for the 5-HTTLPR allele groups with a one-way ANOVA. For attentional engagement, there was not a significant allele group effect,  $F(2, 141) = 0.56, p = .57$  (see Figure 3B). In contrast, for attentional disengagement, there was a significant allele group effect,  $F(2, 141) = 3.79, p = .03$ . Furthermore, group comparisons indicated that the S'S' allele group had significantly greater difficulty disengaging their attention from emotional stimuli than the S'L' and the L'L' groups ( $ps < .01$ ). The S'L' and the L'L' groups did not differ from each other ( $p = .44$ ; see Figure 3C). Mean disengagement scores (and standard errors) collapsing across stimulus valence were LL:  $-7.64 (5.99)$ , SL:  $-2.32 (3.64)$ , SS:  $14.05 (5.27)$ .



**Figure 3.** A: Mean attentional bias, B: attentional engagement, and C: attentional disengagement, with standard errors presented as a function of stimulus valence and serotonin transporter-linked polymorphic region (5-HTTLPR) allele status for Study 2. Note that S'S' includes the SS, L<sub>G</sub>L<sub>G</sub>, and SL<sub>G</sub> allele groups, S'L' includes the SL<sub>A</sub>, L<sub>G</sub>L<sub>A</sub> allele groups, and L'L' includes the L<sub>A</sub>L<sub>A</sub> allele group.

We also examined whether the engagement and disengagement scores were significantly different from zero for each of the genetic groups. For engagement scores, none of the groups were significantly different from zero ( $t_s < |1.5|$ ,  $p_s > .13$ ). However, for disengagement scores, the S'S' group was significantly greater than zero,  $t(28) = 2.66$ ,  $p = .01$ . The S'L' and L'L' groups were not significantly different from zero ( $t_s < |1.2|$ ,  $p_s > .21$ ). People homozygous for the low-expressing 5-HTTLPR allele had particular difficulty disengaging their attention from emotional cues relative to neutral cues.<sup>1</sup>

### General Discussion

Results from these two studies suggest that the low-expressing 5-HTTLPR allele (i.e., S or L<sub>G</sub>) is significantly associated with attentional biases for emotional information among healthy, non-depressed individuals. Furthermore, difficulty disengaging attention from emotional information appeared to be primarily responsible for these effects. More specifically, in Study 1, short 5-HTTLPR allele carriers took significantly longer than long allele homozygotes to disengage their attention from happy and sad stimuli relative to neutral stimuli. In Study 2, individuals homozygous for the low-expressing 5-HTTLPR allele (i.e., S, L<sub>G</sub>) took significantly longer than individuals homozygous for the high-expressing allele (i.e., L<sub>A</sub>) to disengage their attention from happy, sad, and fear stimuli relative to neutral stimuli. These studies are among the first to demonstrate that attentional biases for emotional stimuli may have a genetic component.

Identifying the genetic etiology of attentional biases for emotional information may have important implications for understanding vulnerability to emotional disorders. For instance, the ability to effectively regulate attention away from emotional stimuli has been associated with current depression (Koster et al., 2005), depression vulnerability (Beevers & Carver, 2003), and enhanced emotional reactivity (MacLeod, Rutherford, Campbell, Ebsworthy, & Holker, 2002). Furthermore, Compton (2000) found that difficulty disengaging attention from an invalidly cued location was associated with experiencing heightened negative affect after viewing a distressing film. Future work should examine whether the 5-HTTLPR polymorphism hampers effective emotion regulation via its influence on the disengagement of attention from emotional stimuli. Additional research that measures genotype, biased attention, and emotion regulation is needed to test this possibility.

In contrast to the associations between 5-HTTLPR genotype and difficulty disengaging attention, we did not observe any attentional engagement effects. This is likely due to duration of stimulus presentation. Indeed, as cue duration increases, attention for the location of the cue becomes inhibited in favor of new locations (Posner & Petersen, 1990). A 1,500-ms cue duration used in the present study likely provides sufficient time to orient attention toward the cued location and is therefore not likely to detect enhanced engagement of attention. In our previous work, however, we found significant genotype associations with biased attention for anxiety-related stimuli that were presented for much briefer durations (e.g., 14 and 750 ms; Beevers et al., 2007). Future work should consider including a variety of cue durations to identify how the 5-HTTLPR polymorphism is associated with attentional engagement and disengagement over the time course of attentional

processing. Methodologies that allow for more continuous assessment of attention processing (e.g., eye tracking, evoked response potential) may be particularly useful for clarifying the time course of these effects.

It was intriguing that individuals homozygous for the low-expressing 5-HTTLPR alleles had difficulty disengaging their attention from negative and positive stimuli. It may be that the short 5-HTTLPR allele carriers have increased sensitivity to emotion in general. There is neurological evidence to suggest that this is the case. For instance, Canli et al. (2005) reported that the 5-HTTLPR polymorphism was associated with differential neural activation in response to negative and positive stimuli in limbic, striatal, and cortical regions. They suggested that genetically influenced serotonin transport efficiency may have a greater role in how emotional information in general is processed at a neural level than previously thought. The present findings are consistent with this conclusion and extend the Canli et al. (2005) findings to the psychological level.

Future research examining the 5-HTTLPR polymorphism should also consider simultaneously examining a second 5-HTT polymorphism, the SLC6A4 intron2 VNTR polymorphism. The 10-repeat SLC6A4 intron2 VNTR allele is associated with relatively decreased 5-HTT expression in comparison with the 12-repeat allele (Hranilovic et al., 2004). Thus, if relatively increased 5-HT signaling is partly responsible for the effects observed in this study, then 10-repeat allele carriers should exhibit more biased attention and greater difficulty disengaging attention from emotional stimuli than 12-repeat carriers. Convergence of findings across the 5-HTTLPR and SLC6A4 intron 2 VNTR polymorphisms would provide compelling evidence for the role of 5-HT signaling in biased attention for emotional stimuli. We hope future research will be able to address this issue.

Neuroimaging work may also provide important insights into the neural mediators of difficulty disengaging from emotional material. For instance, there is remarkably consistent evidence for the effect of the 5-HTTLPR polymorphism on amygdala reactivity (for a meta-analysis, see Munafò, Brown, & Hariri, 2008), which may be due to the effect of the 5-HTTLPR polymorphism on downstream targets such as the 5-HT1A autoreceptor (Hariri & Holmes, 2006). Imaging studies with adults suggest that the 5-HTTLPR polymorphism is associated with structural alterations and disrupted functional activation in cortical and limbic areas that are critical for the processing of emotional stimuli (Pezawas et al., 2005). It may be that short 5-HTTLPR allele carriers have poorer prefrontal cortical control over automatic neural responses in the amygdala, which in turn contributes to greater difficulty disengag-

<sup>1</sup> We also conducted analyses using the traditional (or biallelic) 5-HTTLPR polymorphism groups (i.e., not accounting for the L<sub>A</sub> and L<sub>G</sub> variants of the L allele). The Cue Validity (valid, invalid) × Stimulus Valence (sad, happy, fear, neutral) × 5-HTTLPR Allele Type (SS, SL, LL) interaction approached significance for the prediction of latency to identify the target,  $F(6, 420) = 1.98$ ,  $p = .07$ ,  $\eta^2 = .03$ . Furthermore, the Attentional Bias (engagement, disengagement) × Stimulus Valence (sad, happy, fear) × Allele Status (SS, SL, LL) interaction was statistically significant,  $F(2, 140) = 3.77$ ,  $p < .05$ ,  $\eta^2 = .05$ . Results were therefore largely consistent for the biallelic and triallelic 5-HTTLPR classifications in Study 2.

ing their attention from emotional stimuli. Additional work is needed to test this hypothesis.

Several limitations of this study should be noted. Recent research with the spatial cuing task suggests that briefly presented emotional cues may impart greater slowing effects on motor responses than neutral cues. For instance, Mogg et al. (2008) demonstrated significant motor slowing effects on reaction time when emotional cues were presented for 200 ms. Whether similar motor slowing effects are observed for cues that are presented for much longer durations, as in the present study (i.e., 1,500 ms), remains to be determined. Furthermore, it is important to note that the assessment of general biased attention (see Equation 1) in this paradigm is not influenced by motor slowing effects. Nevertheless, future work in this area would be well served to measure the impact of emotional cues on motor responses in order to control for any effects (for an example, see Mogg et al., 2008). Doing so could help to more precisely identify the mechanisms that produce increased reaction time responses to the invalid emotional cues relative to neutral cues among individuals homozygous for the low-expressing 5-HTTLPR polymorphism.

We also examined only a relatively new single nucleotide polymorphism in the long version of the 5-HTTLPR promoter polymorphism (Hu et al., 2005) in Study 2. However, it is encouraging that the results were largely consistent across studies even though we examined only the deletion polymorphism in Study 1. Analyses from Study 2 also indicated that results obtained from the traditional biallelic classification were highly consistent with those using the triallelic classification (see Footnote 1). In addition, the spatial cuing task used in the present study was briefer than versions used in past research, in part because this task was presented in the context of a larger study, and we wanted to minimize participant burden. Future research should consider using spatial cuing task with more trials.

As with any genetic association study, population stratification is a potential concern (Hutchison, Stallings, McGeary, & Bryan, 2004). Population stratification occurs when cases and controls differ with respect to their ethnic background or another variable that may have led to a pattern of nonrandom mating. In our study, this confound is unlikely as 5-HTTLPR allele frequencies did not differ across race or ethnicity in Study 1. Allele frequencies did differ across race in Study 2; however, race was unrelated to attentional bias and statistically controlling for race had no effect on our findings. Third variable explanations, such as the possibility that the 5-HTTLPR promoter polymorphism is in linkage disequilibrium with another functional genetic marker, should also be considered as explanations for the observed effects.

Despite these limitations, we believe this study makes an important and interesting contribution to understanding depression vulnerability. Individuals who inherit two copies of the low-expressing variants of the 5-HTTLPR gene may be more sensitive to life stress and thus at greater risk for depression because they have greater difficulty disengaging their attention from emotional relative to neutral information. Additional work is now needed that examines complex etiological models of depression that link these various mechanisms of risk. By studying mechanisms of risk across levels of analyses (genetic, neural, cognitive, environmental), we may be able to develop more comprehensive models of depression vulnerability and further researchers' understanding of this debilitating disorder.

## References

- American Psychiatric Association. (1994). *Diagnostic and statistical manual of mental disorders* (4th ed.). Washington, DC: Author.
- Beck, A. T., Brown, G. K., & Steer, R. (1996). *Beck Depression Inventory (BDI-II)*. San Antonio, TX: Psychological Corporation.
- Beck, A. T., Steer, R. A., & Garbin, M. G. (1988). Psychometric properties of the Beck Depression Inventory: Twenty-five years of evaluation. *Clinical Psychology Review, 8*, 77–100.
- Beevers, C. G., & Carver, C. S. (2003). Attentional bias and mood persistence as prospective predictors of dysphoria. *Cognitive Therapy and Research, 27*, 619–637.
- Beevers, C. G., Gibb, B. E., McGeary, J. E., & Miller, I. W. (2007). Serotonin transporter genetic variation and biased attention for emotional word stimuli among psychiatric inpatients. *Journal of Abnormal Psychology, 116*, 208–212.
- Bertolino, A., Arciero, G., Rubino, V., Latorre, V., De Candia, M., Mazza, V., et al. (2005). Variation of human amygdala response during threatening stimuli as a function of 5-HTTLPR genotype and personality style. *Biological Psychiatry, 57*, 1517–1525.
- Canli, T., Omura, K., Haas, B. W., Fallgatter, A., Constable, R. T., & Lesch, K. P. (2005). Beyond affect: A role for genetic variation of the serotonin transporter in neural activation during a cognitive attention task. *PNAS, 102*, 12224–12229.
- Caspi, A., Sugden, K., Moffitt, T. E., Taylor, A., Craig, I. W., Harrington, H., et al. (2003, July 18). Influence of life stress on depression: Moderation by a polymorphism in the 5-HTT gene. *Science, 301*, 386–389.
- Compton, R. J. (2000). Ability to disengage attention predicts negative affect. *Cognition & Emotion, 14*, 401–415.
- Cunningham, W. A., Van Bavel, J. J., & Johnsen, I. R. (2008). Affective flexibility: Evaluative processing goals shape amygdala activity. *Psychological Science, 19*, 152–160.
- Don, R. H., Cox, P. T., Wainwright, B. J., Baker, K., & Mattick, J. S. (1991). 'Touchdown' PCR to circumvent spurious priming during gene amplification. *Nucleic Acids Research, 19*, 4008.
- Drevets, W. C. (2000). Neuroimaging studies of mood disorders. *Biological Psychiatry, 48*, 813–829.
- Ekman, P., & Friesen, W. V. (1976). *Pictures of facial affect*. Palo Alto, CA: Consulting Psychologists Press.
- Fallgatter, A. J., Herrmann, M. J., Roemmler, J., Ehlis, A.-C., Wagnen, A., Heidrich, A., et al. (2004). Allelic variation of serotonin transporter function modulates the brain electrical response for error processing. *Neuropsychopharmacology, 29*, 1506–1511.
- Farah, M. J., Wilson, K. D., Drain, M., & Tanaka, J. N. (1998). What is "special" about face perception? *Psychological Review, 105*, 482–498.
- First, M. B., Spitzer, R. L., Gibbon, M., & Williams, J. B. W. (2002). *Structured Clinical Interview for DSM-IV-TR Axis I disorders, research version, patient edition (SCID-IP)*. New York: Biometrics Research, New York State Psychiatric Institute.
- Freeman, B., Powell, J., Ball, D., Hill, L., Graig, I., & Plomin, R. (1997). DNA by mail: An inexpensive and noninvasive method for collecting DNA samples from widely dispersed populations. *Behavioral Genetics, 27*, 251–257.
- Hariri, A. R., Drabant, E. M., Munoz, K. E., Kolachana, B. S., Mattay, V. S., Egan, M. F., et al. (2005). A susceptibility gene for affective disorders and the response of the human amygdala. *Archives of General Psychiatry, 62*, 146–152.
- Hariri, A. R., & Holmes, A. (2006). Genetics of emotional regulation: The role of the serotonin transporter in neural function. *Trends in Cognitive Sciences, 10*, 182–191.
- Hariri, A. R., Mattay, V. S., Tessitore, A., Kolachana, B., Fera, F., Goldman, D., et al. (2002, July 19). Serotonin transporter genetic variation and the response of the human amygdala. *Science, 297*, 400–403.
- Hasler, G., Drevets, W. C., Manji, H. K., & Charney, D. S. (2004).

- Discovering endophenotypes for major depression. *Neuropsychopharmacology*, 29, 1765–1781.
- Hayden, E. P., Dougherty, L. R., Maloney, B., Olino, T. M., Sheikh, H., Durbin, C. E., et al. (2008). Early-emerging cognitive vulnerability to depression and the serotonin transporter promoter region polymorphism. *Journal of Affective Disorders*, 107, 227–230.
- Heils, A., Teufel, A., Petri, S., Stober, G., Riederer, P., Bengel, D., et al. (1996). Allelic variation of the human serotonin transporter gene expression. *Journal of Neurochemistry*, 66, 2621–2624.
- Heinz, A., Braus, D. F., Smolka, M. N., Wrase, J., Puls, I., Hermann, D., et al. (2005). Amygdala-prefrontal coupling depends on a genetic variation of the serotonin transporter. *Nature Neuroscience*, 8, 20–21.
- Heinz, A., Jones, D. W., Mazzanti, C., Goldman, D., Ragan, P., Hommer, D., et al. (2000). A relationship between serotonin transporter genotype and in vivo protein expression and alcohol neurotoxicity. *Biological Psychiatry*, 47, 643–649.
- Hranilovic, D., Stefulj, J., Schwab, S., Borrmann-Hassenbach, M., Albus, M., Jernej, B., et al. (2004). Serotonin transporter promoter and intron 2 polymorphisms: Relationship between allelic variants and gene expression. *Biological Psychiatry*, 55, 1090–1094.
- Hu, X., Oroszi, G., Chun, J., Smith, T. L., Goldman, D., & Schuckit, M. A. (2005). An expanded evaluation of the relationship of four alleles to the level of response to alcohol and the alcoholism risk. *Alcoholism, Clinical and Experimental Research*, 29, 8–16.
- Hutchison, K. E., Stallings, M., McGeary, J., & Bryan, A. (2004). Population stratification in the candidate gene study: Fatal threat or red herring? *Psychological Bulletin*, 130, 66–79.
- Ioannidis, J. P. A., Ntzani, E. E., Trikalinos, T. A., & Contopoulos-Ioannidis, D. G. (2001). Replication validity of genetic association studies. *Nature Genetics*, 29, 306–309.
- Kendler, K. S., Kuhn, J. W., Vittum, J., Prescott, C. A., & Riley, B. (2005). The interaction of stressful life events and a serotonin transporter polymorphism in the prediction of episodes of major depression. *Archives of General Psychiatry*, 62, 529–535.
- Koster, E. H. W., De Raedt, R., Goeleven, E., Franck, E., & Crombez, G. (2005). Mood-congruent attentional bias in dysphoria: Maintained attention to and impaired disengagement from negative information. *Emotion*, 5, 446–455.
- Lecrubier, Y., Sheehan, D. V., Weiller, E., Amorim, P., Bonora, I., Sheehan, K. H., et al. (1997). The Mini International Neuropsychiatric Interview (MINI): A short diagnostic structured interview: Reliability and validity according to the CIDI. *European Psychiatry*, 12, 224–231.
- Lench, N., Stanier, P., & Williamson, R. (1988). Simple non-invasive method to obtain DNA for gene analysis. *Lancet*, 1, 1356–1358.
- Lesch, K. P., Bengel, D., Heils, A., Sabol, S. Z., Greenberg, B. D., Petri, S., et al. (1996). Association of anxiety-related traits with a polymorphism in the serotonin transporter gene regulatory region. *Science*, 274, 1527–1531.
- MacLeod, C., Rutherford, E., Campbell, L., Ebsworthy, G., & Holker, L. (2002). Selective attention and emotional vulnerability: Assessing the causal basis of their association through the experimental manipulation of attentional bias. *Journal of Abnormal Psychology*, 111, 107–123.
- Meulenbelt, I., Droog, S., Trommelen, G. J., Boomsma, D. I., & Slagboom, P. E. (1995). High-yield noninvasive human genomic DNA isolation method for genetic studies in geographically dispersed families and populations. *American Journal of Human Genetics*, 57, 1252–1254.
- Mogg, K., & Bradley, B. P. (1999). Some methodological issues in assessing attentional biases for threatening faces in anxiety: A replication study using a modified version of the probe detection task. *Behavioral Research and Therapy*, 37, 595–604.
- Mogg, K., Holmes, A., Garner, M., & Bradley, B. P. (2008). Effects of threat cues on attentional shifting, disengagement and response slowing in anxious individuals. *Behaviour Research and Therapy*, 46, 656–667.
- Munafò, M. R., Brown, S. M., & Hariri, A. R. (2008). Serotonin transporter (5-HTTLPR) genotype and amygdala activation: A meta-analysis. *Biological Psychiatry*, 63, 852–857.
- Pezawas, L., Meyer-Lindenberg, A., Drabant, E. M., Verchinski, B. A., Munoz, K. E., Kolachana, B. S., et al. (2005). 5-HTTLPR polymorphism impacts human cingulate-amygdala interactions: A genetic susceptibility mechanism for depression. *Nature Neuroscience*, 8, 828–834.
- Posner, M. I. (1980). Orienting of attention. *The Quarterly Journal of Experimental Psychology*, 32, 3–25.
- Posner, M. I., & Petersen, S. E. (1990). The attention system of the human brain. *Annual Review of Neuroscience*, 13, 25–42.
- Posner, M. I., & Rothbart, M. K. (2000). Developing mechanisms of self-regulation. *Development and Psychopathology*, 12, 427–441.
- Roiser, J. P., Cook, L. J., Cooper, J. D., Rubinsztein, D. C., & Sahakian, B. J. (2005). Association of a functional polymorphism in the serotonin transporter gene with abnormal emotional processing in ecstasy users. *American Journal of Psychiatry*, 162, 609–612.
- Sheehan, D. V., Lecrubier, Y., Sheehan, K. H., Amorim, P., Janavs, J., Weiller, E., et al. (1998). The Mini-International Neuropsychiatric Interview (M. I. N. I.): The development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of Clinical Psychiatry*, 59(Suppl. 20), 22–33.
- Spitz, E., Moutier, R., Reed, T., Busnel, M. C., Marchaland, C., Rouberoux, P. L., et al. (1996). Comparative diagnoses of twin zygosity by SSCP variant analysis, questionnaire, and dermatoglyphic analysis. *Behavior Genetics*, 26, 55–64.
- Whalen, P. J. (2007). The uncertainty of it all. *Trends in Cognitive Sciences*, 11, 499–500.
- Wigg, K. G., Takhar, A., Ickowicz, A., Tannock, R., Kennedy, J. L., Pathare, T., et al. (2006). Gene for the serotonin transporter and ADHD: No association with two functional polymorphisms. *American Journal of Medical Genetics Part B, Neuropsychiatric Genetics*, 141, 566–570.
- Zalsman, G., Huang, Y.-y., Oquendo, M. A., Burke, A. K., Hu, X.-z., Brent, D. A., et al. (2006). Association of a triallelic serotonin transporter gene promoter region (5-HTTLPR) polymorphism with stressful life events and severity of depression. *American Journal of Psychiatry*, 163, 1588–1593.

Received March 12, 2008

Revision received March 3, 2009

Accepted March 13, 2009 ■